



- 8. Did you <u>start</u> taking this drug since your last visit?
- 9. [Since your last visit] In what month and year did you start taking this drug?

J	F	M	A	M	J	J	A	S	0	N	D
98	99	00	01	02	03	04	05	06	07	08	09

10. Since your last visit in (MONTH), how long have you used (DRUG)?

One week or less

- O More than 1 week but less than 1 month
- 1-2 months (includes 2 months and longer, but less than 3 months)
- \bigcirc 3–4 months (includes 4 months and longer, but less than 5 months)

◯ 5–6 months

O More than 6 months

11. Did you stop taking this drug, for 2 days or longer, at any time since your last visit? [DOES NOT INCLUDE ALTERNATING DRUG USE]

O NO (GO TO Q13) OYES

12. Why did you stop taking this drug? (MARK ALL THAT APPLY)

- Low white blood cells (low neutrophils)
- Anemia (low red blood cells/low hemoglobin)
- Blood in urine
- OBleeding
- O Dizziness/Headaches
- O Nausea/Vomiting
- Abdominal pain (pancreatitis/abdominal bloating/cramps)
- O Diarrhea
- Muscle pain or weakness (myopathy/myositis/muscle cramps/spasms)
- Burning/tingling in extremities (neuropathy/neuritis/numbness)
- ◯ Kidney stones
- ◯ Kidney failure
- Rash
- O High blood sugar/Diabetes
- O High cholesterol/High triglycerides
- O Painful urination
- O High blood pressure
- O Abnormal changes in body fat
- Vivid nightmares or dreams
- Liver toxicity (abnormal liver function test)
- O Insomnia or problems sleeping

○ Fatigue

- Increased viral load
- Decreased viral load
- Hospitalized
- O Personal decision
- O Prescription changes by physician
- O Too expensive
- Too much bother, inconvenient (ran out/vacation/unable to fill prescription)
- Changed to another drug in order to decrease the number of pills or dosing frequency
- ◯ Study ended
- Other, specify:

1)
2)
3)

13. On average, how often did you take your medication as prescribed?

- 100% of the time
- 95–99% of the time
- 75–94% of the time
- <75% of the time</p>