

# MEN'S ATTITUDE SURVEY


## MARKING INSTRUCTIONS

- Make dark marks that fill the circle completely.
- Make clean erasures.
- Make NO stray marks.
- DO NOT fold this form.
- Fill in one circle for each question.

### INCORRECT MARKS



### CORRECT MARK



ID NUMBER				
	0	0	0	0
1	1	1	1	1
2	2	2	2	2
3	3	3	3	3
4	4	4	4	4
	5	5	5	5
	6	6	6	6
	7	7	7	7
	8	8	8	8
	9	9	9	9

VISIT NO.	
	0 0
	1 1
	2 2
3	3 3
	4 4
	5 5
	6 6
	7 7
	8 8
	9 9

DATE		
MO.	DAY	YEAR
<input type="radio"/> Jan		9 9
<input type="radio"/> Feb		
<input type="radio"/> Mar	0 0	
<input type="radio"/> Apr	1 1	
<input type="radio"/> May	2 2	
<input type="radio"/> June	3 3	
<input type="radio"/> July	4	
<input type="radio"/> Aug	5	
<input type="radio"/> Sept	6	
<input type="radio"/> Oct	7	99
<input type="radio"/> Nov	8	
<input type="radio"/> Dec	9	

## Combination Drug Treatments for HIV

For these questions "**combination drug treatments**" refers to combinations of drugs—including protease inhibitors—used to keep people who are **already infected** with HIV from getting sick with AIDS. In addition, these drugs may also be used by an HIV negative person following a possible exposure to HIV (e.g., through unsafe sex or needle sharing) to try and **prevent** infection with HIV. This is called "**post-exposure prevention**," or **PEP** for short.

- How much have you heard about the use of combination drug treatments to keep people with HIV from getting sick?
  - ☐ nothing
  - ☐ a little
  - ☐ some
  - ☐ quite a bit
  - ☐ a lot
- How many people do you know with HIV who have taken combination drug treatments?
  - ☐ none
  - ☐ 1 or 2
  - ☐ 3 or more
- How much have you heard about using combination drug treatments to try to prevent infection after a possible exposure to HIV (PEP)?
  - ☐ nothing
  - ☐ a little
  - ☐ some
  - ☐ quite a bit
  - ☐ a lot
- How many people do you know who have used combination drug treatments to try to prevent infection after a possible exposure to HIV (PEP)?
  - ☐ none
  - ☐ 1 or 2
  - ☐ 3 or more
- Have you ever used combination drug treatments to try to prevent infection after a possible exposure to HIV (PEP)?
  - ☐ no
  - ☐ yes

CONTINUED ON PAGE 2

PLEASE DO NOT WRITE IN THIS AREA



0933



# Personal Attitudes

**6.** PLEASE MARK ONE NUMBER TO SHOW HOW MUCH YOU AGREE WITH EACH STATEMENT, GIVING YOUR FIRST, "GUT REACTION . . ."

	Do not agree at all				Strongly Agree
1. HIV is no longer the threat it used to be.	①	②	③	④	⑤
2. I am somewhat more willing to take a chance of getting infected or infecting someone else now that combination drug treatments are available.	①	②	③	④	⑤
3. Being drunk or high makes me more comfortable sexually.	①	②	③	④	⑤
4. Because of combination drug treatments for HIV, I am less concerned about becoming HIV+ or infecting someone.	①	②	③	④	⑤
5. I feel tired of always having to monitor my sexual behavior.	①	②	③	④	⑤
6. HIV is now a controllable disease like diabetes.	①	②	③	④	⑤
7. When I am high or drunk I am more likely to do sexual things I usually wouldn't do.	①	②	③	④	⑤
8. I find it difficult to maintain my commitment to safer sex.	①	②	③	④	⑤
9. I like wild "uninhibited" sexual encounters.	①	②	③	④	⑤
10. It would be more difficult for an HIV+ person to infect a partner through unsafe sex if the HIV+ person was taking combination drug treatments.	①	②	③	④	⑤
11. I like new and exciting sexual experiences and sensations.	①	②	③	④	⑤
12. I am a lot less worried about sexual "slipping" now that treatments may be given after unprotected sex.	①	②	③	④	⑤
13. The availability of combination drug treatments for HIV makes me less worried about having unprotected sex.	①	②	③	④	⑤
14. When I am high I find it difficult to stay within my sexual limits.	①	②	③	④	⑤
15. By taking combination drug treatments, an HIV+ person decreases the chance that he/she will infect a partner with HIV.	①	②	③	④	⑤
16. I am less concerned about having unprotected anal sex now that combination drug treatments are available.	①	②	③	④	⑤
17. An HIV+ person whose level of virus in the blood is undetectable is unlikely to transmit HIV to a sex partner.	①	②	③	④	⑤
18. I am interested in trying out new sexual experiences.	①	②	③	④	⑤
19. It takes a lot of effort to keep my sexual behavior safe.	①	②	③	④	⑤
20. When I am high or drunk I am more likely to have sex with people I ordinarily wouldn't have sex with.	①	②	③	④	⑤



# Sexual Behavior, Last 6 Months: Primary Male Partner

IF NO PRIMARY PARTNER, SKIP TO CASUAL/NON-PRIMARY PARTNER SECTION ON PAGE 4.

7. Before or during sex (sexual intercourse) with your primary partner, how often have you during the last six months . . .

	never	rarely	sometimes	about half	most times	nearly all times	every time
A. . . . gotten drunk or "high" on alcohol?	0	1	2	3	4	5	6
B. . . . used poppers?	0	1	2	3	4	5	6
C. . . . used drugs other than alcohol or poppers?	0	1	2	3	4	5	6

- 8A. What is your primary partner's HIV status? ☐ HIV negative  
☐ HIV positive  
☐ I do not know

→ SKIP TO Q9

→ SKIP TO Q9

If your primary partner is HIV+ . . .

- B. Is he currently taking combination treatments? ☐ no ☐ yes ☐ I don't know
- C. What is his viral load? ☐ undetectable ☐ detectable ☐ I don't know

- 9A. Have you disclosed your HIV status to your primary partner? ☐ no  
☐ yes

If you are HIV+ . . .

- B. have you told your primary partner whether you are taking combination treatments? ☐ no ☐ yes
- C. have you told your partner your viral load? ☐ no ☐ yes ☐ I don't know my viral load

- 10A. Have you and your partner discussed rules for sex with other people? ☐ no  
☐ yes

→ SKIP TO Q11

If yes . . . does your agreement include: (Mark all that apply)

- B. ☐ no sex outside the relationship
- C. ☐ no oral sex with another man
- D. ☐ no anal sex with another man
- E. ☐ discussed topic, but no rules established
- F. ☐ anal sex with others, but only with condom
- G. ☐ other, please describe: \_\_\_\_\_

11. Has your primary partner had sex (sexual intercourse) with other men in the last 6 months? ☐ no  
☐ yes  
☐ not sure



# Sexual Behavior, Last 6 Months: Non-Primary Male Partner(s)

IF NO OTHER PARTNERS, STOP HERE.

12. Before or during sex (sexual intercourse) with your non-primary partner(s), how often have you in the last 6 months ...

	never	rarely	sometimes	about half	most times	nearly all times	every time
A... gotten high or "buzzed" on alcohol?	0	1	2	3	4	5	6
B... used poppers?	0	1	2	3	4	5	6
C... used drugs other than alcohol or poppers?	0	1	2	3	4	5	6

13. Communication: How many of your non-primary partner(s) have you ...

	none	some	all
A... disclosed <u>your</u> HIV status to?	0	1	2
B... asked for <u>their</u> HIV status?	0	1	2
C... talked with about condom use or safer sex?	0	1	2

- 14A. Do you know whether any of your non-primary partner(s) are HIV+?

☐ no ☐ yes → How many?

0	10	20	30	40	50	60	70	80	90
0	1	2	3	4	5	6	7	8	9

If you had sex with any HIV+ non-primary partner(s) ...

- B. Did you ever discuss whether they were taking combination drug treatments?

☐ no ☐ yes → How many?

0	10	20	30	40	50	60	70	80	90
0	1	2	3	4	5	6	7	8	9

- C. Did you ever discuss their viral loads?

☐ no ☐ yes

- D. Did you have unprotected anal sex with any HIV+ non-primary partners?

☐ no ☐ yes → How many?

0	10	20	30	40	50	60	70	80	90
0	1	2	3	4	5	6	7	8	9

15. If you are HIV+ ...

How many of your non-primary partners did you ...

	[0%] none	some	all [100%]
A... tell whether <u>you</u> are using combination treatments?	0	1	2
B... discuss <u>your</u> viral loads with?	0	1	2

PLEASE DO NOT WRITE IN THIS AREA

0	1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---	---